


AD-A111 834


LETTERMAN ARMY INST OF RESEARCH PRESIDIO OF SAN FRANC--ETC F/6 6/16  
DOMESTIC SWINE IN PHYSIOLOGICAL RESEARCH. III. BLOOD GAS AND AC--ETC(U)  
FEB 82 J P HANNON, W Y MOORES  
LAIR-113

UNCLASSIFIED

NL

1 OF 1  
AD-A  
111 834





ADA 111 834

6

INSTITUTE REPORT NO. 113

DOMESTIC SWINE IN PHYSIOLOGICAL RESEARCH

III. Blood Gas and Acid-base Values of Arterial and Venous Blood from  
Young Anesthetized Pigs Maintained under Steady-state Ventilatory Conditions


JOHN P. HANNON, PhD

and

WILLIAM Y. MOORES, MD

DIVISION OF COMBAT CASUALTY CARE

FEBRUARY 1982



DTIC  
ELE  
MAR 10 1982  
S H D

LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO CALIFORNIA 94129

82 03 08 128

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

DTIC FILE COPY

Domestic Swine in Physiological Research. III. Blood Gas and  
Acid-base Values of Arterial and Venous Blood from Young Anesthetized  
Pigs Maintained under Steady-state Ventilatory Conditions  
--Hannon and Moores

Reproduction of this document in whole or in part is prohibited except with the permission of the Commander, Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, the Defense Technical Information Center is authorized to reproduce the document for United States Government purposes.

Destroy this report when it is no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

In conducting the research described in this report, the investigation adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Revision of the Guide for Laboratory Animal Facilities and Care, Institute of Laboratory Animal Resources, National Research Council.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360.5)

*John W. Marshall 12/1/66*  
(Signature and date)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER LAIR Institute Report No. 113	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Domestic Swine in Physiological Research. III. Blood Gas and Acid-base Values of Arterial and Venous Blood from Young Anesthetized Pigs Maintained under Steady-state Ventilatory Conditions		5. TYPE OF REPORT & PERIOD COVERED Interim Jun 81 - Dec 81
7. AUTHOR(s) John P. Hannon, PhD and William Y. Moores, MD		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Division of Combat Casualty Care Letterman Army Institute of Research Presidio of San Francisco, CA 94129		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command Fort Detrick, Frederick, MD 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Project 3M161102BS10 Task BA; Work Unit 256
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) US Army Medical Research and Development Command Fort Detrick, Frederick, MD 21701		12. REPORT DATE February 1982
		13. NUMBER OF PAGES 34
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Blood gases; acid-base status; swine; normal values		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The arterial and venous blood gas and acid-base characteristics of anesthetized, young domestic swine were determined under steady-state ventilatory conditions designed to establish arterial P O <sub>2</sub> at 100 torr and pH at 7.40. Under these circumstances, mean (N=15) femoral artery values were: P O <sub>2</sub> , 97 torr; S O <sub>2</sub> , 94%; C O <sub>2</sub> , 15.4 ml/dl; pH, 7.399, P CO <sub>2</sub> , 47 torr; [HCO <sub>3</sub> ], 27.6 mEq/l; and [B.E.], 2.8 mEq/l. Values for pulmonary artery, mixed venous blood were: P O <sub>2</sub> , 36 torr; S aO <sub>2</sub> , 51%.		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

C O<sub>2</sub>, 8.5 ml/dl; pH, 7.335; P CO<sub>2</sub>, 57 torr; [HCO<sub>3</sub>], 29.6 mEq/l; and [B.E.], 3.9 mEq/l. Comparisons of venous values obtained from various vascular sites (pulmonary artery, anterior vena cava, posterior vena cava, internal jugular vein, femoral vein and coronary sinus) revealed numerous between-vessel differences in blood gas and acid-base status. The arterial characteristics of this anesthetized preparation differed from those of the conscious pig; the anesthetized animal had lower pH, [HCO<sub>3</sub>] and [B.E.] values and higher P O<sub>2</sub> and P CO<sub>2</sub> values than the conscious animal. Anesthesia with mechanical ventilation appeared to produce defects in alveolar-arterial gas exchange similar to those reported for other species. The anesthetized pig, nevertheless, can serve as an effective biomedical model for human oriented research.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	
A	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

# ABSTRACT

The arterial and venous blood gas and acid-base characteristics of anesthetized, young domestic swine were determined under steady-state ventilatory conditions designed to establish arterial  $P_{O_2}$  at 100 torr and pH at 7.40. Under these circumstances, mean (N=15) femoral artery values were:  $P_{O_2}$ , 97 torr;  $S_{O_2}$ , 94%;  $C_{O_2}$ , 15.4 ml/dl; pH, 7.399;  $P_{CO_2}$ , 47 torr;  $[HCO_3^-]$ , 27.6 mEq/l; and [B.E.], 2.8 mEq/l. Values for pulmonary artery, mixed venous blood were:  $P_{O_2}$ , 36 torr;  $S_{aO_2}$ , 51%;  $C_{O_2}$ , 8.5 ml/dl; pH, 7.335;  $P_{CO_2}$ , 57 torr;  $[HCO_3^-]$ , 29.6 mEq/l; and [B.E.], 3.9 mEq/l. Comparisons of venous values obtained from various vascular sites (pulmonary artery, anterior vena cava, posterior vena cava, internal jugular vein, femoral vein and coronary sinus) revealed numerous between-vessel differences in blood gas and acid-base status. The arterial characteristics of this anesthetized preparation differed from those of the conscious pig; the anesthetized animal had lower pH,  $[HCO_3^-]$  and [B.E.] values and higher  $P_{O_2}$  and  $P_{CO_2}$  values than the conscious animal. Anesthesia with mechanical ventilation appeared to produce defects in alveolar-arterial gas exchange similar to those reported for other species. The anesthetized pig, nevertheless, can serve as an effective biomedical model for human oriented research.

## PREFACE

This is the third in a series of studies concerned with the utility of domestic swine as a large animal biomedical model for human-oriented physiological experiments. Earlier reports provided a brief review of the use of pigs in medical research and a description of the cation and anion characteristics of porcine arterial blood. Future reports will be concerned with the surgical preparation of swine for experiments to be conducted under conscious conditions and with the population characteristics for the blood gas and acid-base status of arterial blood obtained under basal conditions.

We would like to express our appreciation to Virginia L. Gildengorin, Ph.D., for the invaluable assistance she provided in the statistical evaluation of the data reported here, to Lottie B. Applewhite for the many editorial and format improvements incorporated in this report, and especially to Linda Pukajlo for the innumerable hours she spent typing, proofreading, and assembling the manuscript.



## TABLE OF CONTENTS

Abstract . . . . .	i
Preface . . . . .	ii
Table of Contents . . . . .	iii
BODY OF REPORT	
INTRODUCTION . . . . .	1
METHODS . . . . .	2
RESULTS . . . . .	3
DISCUSSION . . . . .	4
CONCLUSIONS . . . . .	9
RECOMMENDATIONS . . . . .	10
REFERENCES . . . . .	11
Appendix (Tables 1-11) . . . . .	15
Distribution List . . . . .	28

**BLANK PAGE**

DOMESTIC SWINE IN PHYSIOLOGICAL RESEARCH. III. Blood Gas and Acid-base Values of Arterial and Venous Blood from Young Anesthetized Pigs Maintained under Steady-state Ventilatory Conditions

Historically, the mongrel dog has served as the predominant large animal model for experimental studies of mammalian cardiovascular function. Such predominance continues at the present time, but with increasing frequency one sees miniature or domestic pigs being used in the conduct of these studies. In most instances, the selection of swine as a biomedical model stems from both anatomical and functional considerations. The anatomy of the coronary vasculature in the pig, for example, more closely resembles that of man than does the coronary vasculature of the dog (1,2). As a consequence, the pig is considered by some to be a superior animal model for studies of coronary blood flow (3,4) and myocardial function (4-7). The domestic pig would also appear to be superior to the dog for investigations of circulatory shock. Hepatic venous outflow in the dog, for example, is regulated by a highly developed sphincter-like mechanism (8) which is quite sensitive to the vasoactive materials produced during shock (9). Closure of this sphincter mechanism in the shocked dog results in hepatic congestion, increased portal vein pressure, and intestinal hemorrhage (9). These responses are rarely seen in humans or swine (9, 10) probably because the sphincter mechanism is poorly developed (11).

As indicated in earlier reports of this series (12, 13), effective use of the domestic pig as a biomedical model is oftentimes encumbered by a lack of readily available experimental data describing the population characteristics for certain physiologic and biochemical variables. To address one such problem, we measured the major arterial anion and cation concentrations in young anesthetized domestic swine maintained under steady state ventilatory conditions, and these values were compared to those published in the literature (13). A paucity of data was encountered also when we attempted to ascertain normal values for the blood gas and acid-base status of these animals. Most values reported in the literature were measured in venous blood samples with little or no attention given

to the physiologic status of the pigs at the time of blood collection. In view of the many factors (eg. ventilation, kidney function, sampling site) that can affect blood gas and acid-base status, it was perhaps not too surprising to find wide divergence in the values obtained by various investigators.

The study reported here had two objectives. The first was to determine whether or not the anesthetized domestic pig could be maintained under steady state ventilatory conditions such that arterial pH and  $PO_2$  were established at levels commonly seen in humans, that is, about 7.40 and 100 torr respectively. The second objective was to provide, in the same animal model, data on the blood gas and arterial and venous samples obtained from vascular sites that are commonly monitored in cardiovascular research studies.

#### METHODS

Fifteen young domestic pigs (Hampshire - Duroc Cross) of both sexes, 60 to 90 days old, were used. They were obtained from a commercial breeder (J.G. Boswell, Corcoran CA) and, upon receipt at the laboratory, were maintained in a common indoor holding area until utilized in the study, usually within two weeks after arrival. They were fed a commercial ration (Purina Pig Chow, Ralston-Purina Co. St. Louis, MO) and received water ad libitum.

After an overnight fast, each pig was brought into the laboratory and anesthetized. Halothane was used for induction, meperidine hydrochloride and nitrous oxide for maintenance, and succinylcholine for paralysis. The animal was placed in a supine position, and ventilation was provided by means of an endotracheal tube and a mechanical ventilator (Ohio Medical Products, Madison, WI). The latter was used also to regulate nitrous oxide and oxygen delivery. Ventilation and oxygen delivery were adjusted to establish an arterial  $P O_2$  of approximately 100 torr and a pH of approximately 7.40. This was ascertained by periodic sampling of femoral artery blood obtained by means of an indwelling catheter. No infusions of acid or base were required to maintain steady-state acidity levels. Normothermia (37C) was maintained by means of a circulating warm water heating pad. The pig remained under these conditions for approximately one hour, during which a sternotomy was performed and indwelling catheters were placed in the femoral vein, anterior and posterior vena cava, pulmonary artery, internal jugular vein, and coronary sinus.

When  $P O_2$  and pH had been stabilized for at least one-half hour without any adjustments of pulmonary minute volume or inspired  $O_2$  tension, 3 ml samples of blood were drawn from each of the catheters for blood gas and acid-base determinations. The total time involved

in obtaining these samples was 30 seconds or less. Heparin (1000 units/ml) was used to fill the dead space within the syringes, and care was exercised in avoiding sample contamination with air bubbles. Immediately after withdrawal the syringes were capped and immersed in ice water where they remained until the blood gas and acid-base measurements were completed. These measurements were made with an automated blood gas analyzer (Model 813, Instrumentation Laboratory, Lexington, MA), which had been calibrated with precision buffers and analyzed gases provided by the manufacturer. In addition, hemoglobin (Hb) concentration was determined with a cooximeter (Instrumentation Laboratory, Lexington, MA), and the oxygen saturation ( $S_{O_2}$ ) of all samples was measured with an oximeter (American Optical Co., Buffalo, NY). Oxygen content (ml/dl of blood) was calculated as the product of  $1.34 \text{ (ml } O_2/\text{gHb)} \times \text{Hb (g/dl)} \times SO_2 (\%) \times 10^{-2}$ . All data were summarized in terms of the mean population value, standard deviation, standard error of the mean, and range. The significance of differences in venous values were evaluated with single-factor analyses of variance. Interrelationships between variables were evaluated with Pearson Product-Moment correlation values.

#### RESULTS

#### RESULTS

Body weight, hemoglobin, and ventilatory functions recorded at the time of blood sampling are summarized in Table 1. Tables 2 through 8 summarize the blood gas and acid-base data of samples collected from various arterial and venous sites. These data show that reasonable success was achieved in establishing a ventilatory steady state in which the arterial (femoral) pH was maintained at about 7.40 and  $P_{O_2}$  at about 100 torr.

In viewing these tables, marked differences in the blood gas and acid-base characteristics of arterial and venous blood, including mixed-venous blood from the pulmonary artery, are readily evident. Not so readily evident are the differences between venous blood samples. Analyses of variance evaluation thus revealed significant between-vessel differences for all of the venous blood gas and acid-base variables except bicarbonate (Tables 9a, 9b). In general, more between-vessel differences were found for oxygen variables than for  $CO_2$  or other acid-base variables.

Table 10 summarizes the possible interrelationships between key blood gas and acid-base variables for arterial and mixed venous blood. Significant correlation coefficients for certain of these interrelationships, such as pH,  $P_{CO_2}$ ,  $[HCO_3^-]$  and [B.E.] might be anticipated, since they all reflect characteristics of a common functional entity. Others, however, were unexpected. The latter included the significant positive correlation coefficients that were obtained between  $P_{O_2}$  and the  $P_{CO_2}$ , bicarbonate

concentration and base excess concentration of blood taken from the femoral artery; these correlations were not significant in mixed venous blood taken from the pulmonary artery.

Possible correlations between ventilatory variables and key arterial blood gas and acid-base variables are summarized in Table 11. A highly significant positive correlation was found between arterial  $P O_2$  and the percentage oxygen in the inspired air ( $F_I O_2$ ), an effect that would be expected. But unexpectedly, a highly positive correlation was found between  $P CO_2$  and  $F_I O_2$  and, albeit of borderline significance, between  $F_I O_2$  and bicarbonate concentration. Respiratory frequency also showed a positive correlation of borderline significance to arterial  $P CO_2$ . Inspiratory minute volume and tidal volume showed no significant relationships to blood gas and acid-base status, although the negative correlation between  $P CO_2$  and tidal volume approached statistical significance.

## DISCUSSION

The results of this study show that it is indeed possible to establish steady state ventilatory conditions in which the values for arterial pH and  $P O_2$  of anesthetized swine simulate those of the normal human. The recorded data reveal, however, two potential problems that must be addressed if the present simulation model is used for human-oriented investigations. One of these was the high porcine values, relative to those characteristic of humans (14, 15) for arterial and venous  $P CO_2$ , bicarbonate, and base excess. The other problem concerned potential distortion of normal alveolar-pulmonary capillary gas exchange by the mechanical ventilation procedures used to establish steady state values for pH and  $P O_2$ .

Were the elevated acid-base values recorded in this study truly characteristic of the normal pig or were they the result of the experimental conditions imposed on the animal before the blood gas and acid-base measurements were made? This question can not be addressed by the experimental work reported here, and data reported in the literature, for the most part, fail to resolve the issue. Reported porcine venous values for pH are higher (16-24), lower (18, 25-27), or about the same (18, 20, 25, 26, 28-30) as those obtained in the present study. Most reported values for venous  $P CO_2$  are lower than those recorded in the present study (16-20, 22, 24, 26-30), but frequently equivalent values are seen (18, 20, 23, 25). Venous bicarbonate and base excess concentrations reported in the literature, relative to those reported here, show about the same variability as  $P CO_2$ .

On the arterial side, far fewer blood gas and acid-base measurements have been reported. In general, it would appear that most of

the reported pH values (18, 21, 23, 28) are higher than those recorded here, but occasional exceptions are seen (31). In contrast, most reported  $P_{CO_2}$  values (18, 25, 28, 31), again with occasional exceptions (18), are lower than those recorded here. Measurements of bicarbonate and base excess concentrations have been so few and the reported data so divergent (18, 23, 28) that it is nearly impossible to discern characteristics of the normal pig from published values.

Inconsistencies in the published values for the blood gas and acid-base characteristics of the normal pig can be attributed to a variety of factors. Foremost among these were disparities in the physiological status of the animal at the time blood samples were taken for analysis. In almost every instance, control or supposedly normal values were obtained as a prelude to some experimental manipulation (e.g. exercise, heat stress, shock) of immediate interest to the investigator. In most instances, there is little evidence that steady-state metabolic and ventilatory conditions were established before these control measurements were made. Local or general anesthetics, or physical restraint were oftentimes employed to secure the requisite blood samples for study. These manipulations, obviously, can readily alter the physiological status of the animal and consequently the outcome of blood gas and acid-base measurements. Other variables include differences in animal size, environmental temperature at the time of study, animal breed, and, in the case of venous blood, the vascular sampling site. The distinctly different blood gas and acid-base characteristics recorded here for two commonly used veins, the jugular and the femoral, illustrate this last point.

Ideally, control or normal values should approximate those obtained under one widely-recognized standard condition. The basal metabolic state (i.e. the metabolic state characteristic of a recumbent, well-rested, conscious subject who has been fasted overnight) represents, perhaps, the most common reference standard. Few attempts have been made to acquire data from pigs in this condition. This is understandable since most pigs are excitable animals, and some sort of physical or chemical restraint is usually required before experimental measurements can be made. One way to minimize, or eliminate, the restraint variable is to obtain blood samples from the conscious animal by means of chronically implanted vascular catheters. A few investigators have reported blood gas and acid-base values for samples so acquired (18, 21, 23, 28), but, unfortunately, their reports contain little information about the metabolic state of the animals at the time of study.

Basal conditions were approached, if not achieved, in a recent investigation of hemorrhagic hypotension in conscious pigs (32). In this study, the animals received chronically-implanted arterial catheters 7 to 10 days before they were subjected to experimentation.

At the time control samples were taken for blood gas and acid-base evaluation, they had been fasted overnight and had been in a conscious, unrestrained, recumbent position for 30 to 60 minutes before any measurements were made. Under these conditions, 18 pigs had an average pH of 7.50, a  $P_{CO_2}$  of 40.3 torr, a  $P_{O_2}$  of 81.6 torr, bicarbonate concentration of 30.4 mEq/l, and a base excess concentration of 7.5 mEq/l. An untreated control group (N=6) included in this study showed little deviation from the foregoing values when the measurements were repeated at hourly intervals over a 6-hour period.

All of the blood gas and acid-base values obtained in the present investigation differ appreciably from values previously recorded in conscious pigs. In large measure these differences appear attributable to the establishment of a steady-state arterial pH at 7.40. This resulted in  $P_{CO_2}$  values that were higher and  $[HCO_3^-]$  values that were lower than those observed in the conscious animal. The bicarbonate difference, furthermore, appeared responsible, at least in part, for the lower base excess values reported here, between the two studies the arterial bicarbonate difference was 2.8 mEq/l, the base excess difference 5.7 mEq/l. Additionally, arterial  $P_{O_2}$  in the conscious animal is considerably lower than the 100 torr steady-state value selected for the present study. The value in the conscious pig, furthermore, was lower than the value usually seen in the conscious human (14, 33). Arterial oxygen saturation, it might be noted, is lower also because of the rightward position of the porcine oxyhemoglobin dissociation curve relative to that of man (34).

In view of the foregoing, it is clearly evident that researchers using anesthetized swine as a biomedical model must weigh the merits of establishing steady-state blood gas and acid-base conditions which simulate those of humans against those which simulate the normal conscious pig.

The second potential problem raised by this study may have more serious consequences if the biomedical model described here is to be used for routine cardiovascular or other studies. The problem centers on the oxygen fraction of inspired air that was needed to maintain arterial  $P_{O_2}$  at the desired level (Table 1), and the positive correlation of arterial  $P_{O_2}$  and  $P_{CO_2}$  that was obtained once steady state conditions were established. Taken together, these observations suggest that the selected ventilatory and arterial parameters utilized in this study may lead to an impairment of normal alveolar-arterial gas exchange.

Support for this implication is found when the alveolar to arterial (A-a) oxygen gradient of these pigs is compared to those of conscious humans or conscious pigs. Since alveolar oxygen tension ( $P_{AO_2}$ ) values were not obtained in the present study nor,



insofar as can be determined, in other studies of swine, these comparisons must be based on estimated values. Accordingly, if certain assumptions are made,  $P_{A}O_2$  can be calculated from the alveolar gas equation (see Otis (35) for derivation):

$$P_{A}O_2 = P_{I}O_2 - \frac{P_{A}CO_2}{F_{I}O_2} + F_{I}O_2 \frac{P_{A}CO_2}{R} - P_{A}CO_2$$

For present circumstances, it will be assumed that  $F_{I}O_2 = 0.209$  and  $P_{I}O_2 = 149$  torr for conscious humans and pigs breathing air at sea level. The average steady-state  $F_{I}O_2$  for the anesthetized pigs in this study was 0.380 which would be equivalent to a  $P_{I}O_2$  value of 271 torr. Since  $CO_2$  has a high membrane diffusion capacity (36) equality of  $P_{A}CO_2$  and  $P_{a}CO_2$  will be assumed (i.e. 40 torr for conscious pigs and humans and 47 torr for anesthetized pigs). Finally, an R value of 0.80 will be assumed for all conditions, and measured (32) or commonly reported (14) values for  $P_{a}O_2$  will be used in the calculations of  $P_{A}O_2$ . The alveolar equation thus implemented yields the following characteristics for swine and humans:

	$P_{A}O_2$	$P_{a}O_2$	$P_{A}O_2 - P_{a}O_2$
Conscious human	101	90	11
Conscious pig	101	82	19
Anesthetized pig	217	97	120

On the basis of this tabulation it is readily evident that the A-a gradient of the anesthetized pigs used in this study was approximately 6 times that of the conscious pig and 10 times that of the conscious human. The accuracy of these A-a gradient estimations, obviously, hinges on the accuracy of the various assumed values which were inserted into the alveolar gas equation. Of these, the most likely deviation would be the assumed  $P_{A}CO_2$  values; they may have been considerably lower than  $P_{a}CO_2$  values. A lower  $P_{A}CO_2$  value, however, would lead a higher calculated value for  $P_{A}O_2$  and, consequently, a higher A-a gradient. Thus, if  $P_{A}CO_2$  were 40 torr in anesthetized pigs, the calculated  $P_{A}O_2$  would be 225 torr and the A-a gradient 128 torr.

The factors responsible for the positive correlation between steady-state values for  $P_{aO_2}$  and  $P_{aCO_2}$  seen here (Table 10) were not investigated. Under normal circumstances one would expect a negative correlation between the two variables. For example, if ventilation were increased by voluntary or mechanical means then  $P_{aO_2}$  should increase also while  $P_{aCO_2}$  should decrease. The existence of a positive correlation, therefore, suggests the presence of an impediment to normal alveolar-arterial exchange of both oxygen and carbon dioxide. In this regard, it is important to note that a positive correlation was not obtained for venous blood taken from the pulmonary artery, or for that matter for blood from any other venous site. The effect, therefore, arose during the transit of blood through the lungs.

In the establishment of steady-state values for  $pH_a$  and  $P_{aO_2}$ , both  $F_{IO_2}$  and ventilatory minute volume were manipulated. If preliminary measurements showed  $P_{aO_2}$  values to be too low then the  $F_{IO_2}$  of the inspired air was increased. The opposite maneuver was used if  $P_{aO_2}$  values were too high. Similarly, if preliminary  $pH_a$  values were too low (or high) ventilation would be increased (or decreased) to enhance (or attenuate) carbon dioxide elimination and so shift arterial pH values in the desired direction. After steady state conditions had been established, the selected fraction of oxygen in the inspired air had a much greater impact on arterial blood gas and acid-base status than the selected level of ventilation. Thus,  $P_{aO_2}$  showed, as would be anticipated, a positive correlation to  $F_{IO_2}$  (Table 11). But unexpectedly,  $F_{IO_2}$  also showed a highly positive correlation with  $P_{aCO_2}$ . Under steady-state conditions one would expect these two variables to be unrelated. One might anticipate also a negative correlation between ventilatory minute volume or tidal volume and  $P_{aCO_2}$ . The latter was not seen, however (Table 11).

What could account for these apparently paradoxical results? An impediment to normal alveolar-pulmonary capillary gas exchange, incurred during the establishment of the steady state, would seem to be one possibility. If such an impediment occurred shortly after the animal was placed on the mechanical ventilator, then  $P_{aO_2}$  would tend to decrease while  $P_{aCO_2}$  would tend to increase. The increase in  $P_{aCO_2}$  would be a desired effect since it would be required in the establishment of steady-state  $pH_a$  at 7.40. That is, if the  $pH_a$  was about 7.50 when the pig was conscious (32), then a  $P_{aCO_2}$  increase would be needed to lower the value to 7.40 when it was anesthetized. In most instances, defects in alveolar-arterial gas exchange should have a much greater impact on  $P_{aO_2}$  than on  $P_{aCO_2}$ , because  $CO_2$ , compared to  $O_2$ , has a 20-fold greater alveolar membrane solubility and proportionately greater membrane diffusion capacity (36). In the present study, therefore, sizable increases in  $F_{IO_2}$  level were required to achieve the desired steady-state

level for  $P_{aO_2}$ . These increments, it might be assumed, should have had no effect on  $P_{aCO_2}$  since the latter was regulated by ventilatory changes. This assumption, however, may not be valid since Larson and Severinghaus (37) showed in man that increases in  $F_{IO_2}$  led to increases in the a-A gradient for  $CO_2$ . They attributed the gradient increase to an oxygen-induced relaxation of the pulmonary vascular bed which diverted blood flow to the more poorly ventilated portions of the lung. Such a mechanism, if operative in the anesthetized pig, could account for the positive correlation of  $P_{aO_2}$  and  $P_{aCO_2}$  seen in the present study. Accordingly, greater increments in  $F_{IO_2}$  would produce proportionately greater increments in  $P_{aO_2}$  (Table 11), but they would also produce proportionately greater increments in the a-A gradient for  $CO_2$  and proportionately greater increments in  $P_{aCO_2}$  (Table 11).

Defects in pulmonary gas exchange during anesthesia have been widely recognized. In general, they have been attributed to an uneven distribution of alveolar ventilation in relation to alveolar volume and blood flow (38). Anesthetics, including nitrous oxide, cause atelectasis (39), a decrease in pulmonary compliance (40), an increase in dead space (41), venous admixture to arterial blood in the lung with a lowering of  $P_{aO_2}$  (39) and an increase in the a-A gradient for  $CO_2$  (41). It appears that most of these effects can be attributed to an unchanging tidal volume and decreased residual functional capacity, since equivalent defects have been recorded in conscious, mechanically ventilated humans (42).

The problems with pulmonary gas exchange encountered in the present study, therefore, would not seem to be unique to swine and should not seriously detract from their use in biomedical studies requiring anesthetized animals. In fact, this same preparation has been used successfully for extracorporeal circulation experiments lasting several hours (4).

#### CONCLUSIONS

- ° With proper regulation of ventilation and inspired oxygen tension, the arterial  $P_{O_2}$  and pH of anesthetized swine can be brought to steady-state values that approximate those of humans.
- ° Under these circumstances the values for arterial  $P_{CO_2}$ ,  $[HCO_3^-]$  and  $P_{O_2}$  are higher than those seen in conscious swine. Base excess concentration and pH values are lower than those seen in conscious swine.

- ° Under steady-state ventilatory conditions there are significant, between-vessel, differences in the blood gas and acid-base characteristics of venous blood.
- ° Mechanical ventilation in the anesthetized pig leads to impairment of normal alveolar-arterial gas exchange.

#### RECOMMENDATIONS

- ° The feasibility of establishing blood gas and acid-base characteristics of the anesthetized pig at values approximating those seen in the conscious pig should be investigated.
- ° In anesthetized mechanically ventilated swine, attempts should be made to minimize or eliminate defects in alveolar-arterial gas exchange by the inclusion of periodic hyperinflation.

# REFERENCES

1. LUMB, G.D. Experimentally induced cardiac failure in swine. Pathological changes. In: Swine in Biomedical Research, edited by L.K. Bustad and R.O. McClellan. Seattle: Frayn Printing, 1966. pp 389-408
2. GREGG, D.D. and R.E. PATTERSON. Functional importance of the coronary collaterals. N Engl J Med 303: 1403-1406, 1981
3. ENGLEMAN, R.M., S. ADLER, T.H. GAUGE, R. CHANDRA, A.D. BOYD, and L.G. BAUMAN. The effect of normothermic arrest and ventricular fibrillation on coronary flow distribution in the pig. J Thorac Cardiovasc Surg 29: 858-869, 1975
4. MOORES, W.Y., J.P. HANNON, J. CRUM, D. WILFORD, W.G. RODKEY, and J.W. GEASLING. Coronary flow distribution and dynamics during continuous and pulsatile extracorporeal circulation in the pig. Ann Thorac Surg 24: 582-590, 1977
5. IKRAM, H., J.C. NICHOLS, and D.C. FLEMMING. An experimental model for the in vivo study of coronary artery thrombosis. Angiology 26: 356-364, 1975
6. LIEDKE, A.J., H.C. HUGHES, and J.R. NEELY. An experimental model for studying myocardial ischemia. Correlation of hemodynamic performance and metabolism in the working swine heart. J Thorac Cardiovasc Surg 69: 203-211, 1975
7. LEE, K.T. Production of advanced coronary atherosclerosis, myocardial infarction and sudden death in swine. Exp Mol Pathol 15: 170-190, 1971
8. KNISELY, M.H., F. HARDING, and H. DEBACKER. Hepatic sphincters. Brief summary of present-day knowledge. Science 125: 1023-1026, 1957
9. ZWEIFACH, B.W. Aspects of comparative physiology of laboratory animals relative to the problem of experimental shock. Fed Proc 20 (Part III, Suppl 9): 18-29, 1961
10. LINDBERG, B. and N. DARLE. The effect of dextran 40 and blood transfusion on hepatic circulation and oxygen consumption in hemorrhagic shock. J Surg Res 23: 264-273, 1977
11. ELIAS, H. Liver morphology. Biol Rev 30: 263-310, 1955
12. HANNON, J.P. Domestic swine in physiological research. I. A biomedical model. Institute Report No. 91. Presidio of San Francisco: Letterman Army Institute of Research, 1981

13. HANNON, J.P., J.H. SKALA and W.Y. MOORES. Domestic swine in physiological research. II. Electrolyte values for arterial serum from young anesthetized pigs maintained under steady-state ventilatory conditions. Institute Report No. 92. Presidio of San Francisco: Letterman Army Institute of Research, 1981
14. SEVERINGHAUS, J.W. Blood gas concentrations. In: Handbook of Physiology. Respiration, (Vol II, Sect. 3, Chapter 61). Washington, DC: American Physiological Society, 1965, pp 1475-1487
15. SIGGAARD-ANDERSEN, O. The Acid-base Status of the Blood. Baltimore: Williams and Wilkins, 1963
16. BÜNGER, U., N. STEINHARDT, G. RIEHM, and L. LYHS. Änderungen des Milchsäuregehaltes und des Säure-Basen-Status im Blut von Schweinen bei standardisierter submaximaler Laufbelastung. Monatsch Veterinarmed 30: 59-63, 1975
17. LÖWE, G., M. STEINHARDT, and L. LYHS. Untersuchungen zum Säuren-Basen-Status des Blutes und pH - Wert des Muskels bei aufeinander-folgenden Belastungen des Hausschweines. Monatsch Veterinarmed 31: 164-171, 1976
18. VAN DEN HENDE, C., E. MUYLLE, and W. OYAERT. Oxygen utilization and metabolic acidosis after exercise in pigs. Ztbl Veterinarmed Reihe A. 17: 167-173, 1970
19. HARRIS, W.H. Hemoglobin, blood gases and serum electrolyte values in swine. Can Vet J 15: 282-285, 1974
20. FORREST, J.C., J.A. WILL, G.R. SCHNIDT, M.D. JUDGE, and E.J. BRISKEY. Homeostatis in animals (Sus domesticus) during exposure to a warm environment. J Appl Physiol 24: 33-39, 1968
21. TOPEL, D.G., D.E. GALLOWAY, J.A. WILL, N.E. WEIRICH, R.H. GRUNNER, R.G. CASSENS, R.G. KAUFFMAN, and E.J. BRISKEY. Effect of environmental temperature on physiological characteristics of pigs with fast and slow glycolyzing muscle. J Anim Sci 32: 1103-1106, 1971
22. STEINHARDT, M., U. BÜNGER, G. RIEHM, and L. LYHS. Einfluss einer Anämie auf einige Reactionen des Hausschweines bei motorischer Belastung. I. Ausgangs werte und Charakterisierung der Anämie. Arch Exp Vet Med 29: 563-570, 1975
23. SCOTT, D. and G.H. McINTOSH. Changes in blood composition and urinary mineral excretion in the pig in response to acute acid-base disturbance. Q J Exp Physiol 60: 131-140, 1975

24. BARTKO, P., L. VRZGULA, and M. CHYLA. Dynamika indexov acidobázickej homeostázy krvi prasiat do odstavu. *Vet Casopis* 16: 131-137, 1974
25. BARTKO, P. A. MICHNA, M. CHYLA, and L. VRZGULA. Vplyv fyzického zaťaženia na hodnoty indexov acidobázickej homeostázy krvi ošipaných. *Vet Med (Praha)* 20: 339-346, 1975
26. ŠTROS, K., J. KOMAREK, and L. JADRNÝ. Acidobazické hodnoty krve prasete. *Vet Med (Praha)* 20: 327-337, 1975
27. KOMAR, E. Badania nad wpływem znieczulenia ogólnego u świń, wywołanego dożylnym podaniem wodnika chloralu i eunarkonu, na równowagę kwasowo zasadowa we krwi i zmian w poziomie niektórych elektrolitów w surowicy. *Ann Univ Mariae Curie-Skłodowska, Sectio DD* 25: 211-229, 1970
28. JUDGE, M.D., G. EIKELBOOM, Z. ZUIDAM, and W. SYBESMA. Blood acid-base status and oxygen binding during stress-induced hyperthermia in pigs. *J Anim Sci* 37: 776-784, 1973
29. KOMÁREK, K. ŠTROS, I. SÝKORA, I. KYNCLOVA, L. JADRNÝ, and P. SELINGER. Hodnoty acidobazické rovnováhy krve některých druhů domácích zvířat. *Vet Med (Praha)* 21: 51-59, 1976
30. LEBEDA, M. Normální acidobazické hodnoty venózní krve krav a hodnocení stupně kompenzace metabolické poruchy. *Veterinářství* 22: 454-456, 1972
31. SCHÜLKE, B., L. LYHS, and O. KÄDING. Über den Einfluss exogener Hyperthermie und thermischer Hyperventilation auf den Partialdruck des CO<sub>2</sub> im Blut des Schweines. *Arch Exp Vet Med* 22: 633-637, 1968
32. HANNON, J.P., P.B. JENNINGS, Jr. and R.S. DIXON. Physiologic aspects of porcine hemorrhage. IV. Blood gas and acid-base status of the conscious animal following 30 and 50 percent blood volume loss. Institute Report No. 111. Presidio of San Francisco; Letterman Army Institute of Research, 1981
33. ALTMAN, P.I., and D.S. DITMAR. Acid-base balance of blood and plasma: man. Part 1, Acid-base and blood gas values for various ages. In: *Biology Data Book*. (Second edition, Vol III, Sect.252) Bethesda: Federation of American Societies for Experimental Biology, 1974, pp 1830-1832
34. BARTELS, H. and H. HARMS. Sauerstoffdissoziationskurven des Blutes von Säugetieren (Mensch, Kaninchen, Meerschweinchen, Hund, Katze, Schwein, Rind, und Schaft). *Pflügers Arch* 268: 334-365, 1959

35. OTIS, A.B. Quantitative relationships in steady-state gas exchange. In: Handbook of Physiology. Respiration, (Vol I, Sect. 3, Chapter 27). Washington, DC: American Physiological Society, 1964. pp 681-698
36. FORSTER, R.E. Diffusion of gases. In: Handbook of Physiology. Respiration. (Vol I, Sect 3, Chapter 33) Washington, DC: Am Physiol Soc 1964. pp 839-872
37. LARSON, C.P. Jr., and J.W. SEVERINGHAUS. Postural variations in dead space and CO<sub>2</sub> gradients breathing air and O<sub>2</sub>. J Appl Physiol 17: 417-420, 1962
38. FORSTER, R.E. Interpretation of measurements of pulmonary diffusion capacity, In: Handbook of Physiology. Respiration. (Vol II, Sect 3, Chapter 59,) Washington, DC: American Physiological Society 1965, pp 1453-1468
39. BENDIXEN, H.H., J. HEDLEY-WHITE, and M.B. LAVER. Impaired oxygenation in surgical patients during general anesthesia with controlled ventilation: A concept of atelectasis. N Engl J Med 269: 991-996, 1963
40. MEAD, J., and C. COLLIER. Relation of volume history of lungs to respiratory mechanics in anesthetized dogs. J Appl Physiol 14: 669-678, 1959
41. ASKROG, V.F., J.W. PENDER, T.C. SMITH, and J.E. ECKENHOFF. Changes in respiratory dead space during halothane, cyclopropane and nitrous oxide. Anesthesiology 25: 342-352, 1964
42. CAMPBELL, E.J., M.J.F. NUNN, and B.W. PECKETT. A comparison of artificial ventilation and spontaneous respiration with particular reference to ventilation-blood flow relationships. Br J Anaesth 30: 166-175, 1958



#### LEGENDS OF TABLES

- TABLE 1 - Animal Characteristics and Experimental Conditions at the Time of Blood Gas and Acid-Base Measurement
- TABLE 2 - Arterial and Venous P O<sub>2</sub> Values for Young Swine
- TABLE 3 - Arterial and Venous O<sub>2</sub> Saturation Values for Young Swine
- TABLE 4 - Arterial and Venous O<sub>2</sub> Content Values for Young Swine
- TABLE 5 - Arterial and Venous pH Values for Young Swine
- TABLE 6 - Arterial and Venous P CO<sub>2</sub> Values of Young Swine
- TABLE 7 - Arterial and Venous Bicarbonate Values for Young Swine
- TABLE 8 - Arterial and Venous Base Excess Values for Young Swine
- TABLE 9a - Analyses of Variance Evaluation of Differences in Blood Gas and Acid-Base Characteristics of Venous Blood Taken from Various Vascular Sites of Young Pigs
- TABLE 9b - Newman-Keuls Evaluation of Differences in Blood Gas and Acid-Base Characteristics of Venous Blood Taken from Various Vascular Sites of Young Pigs
- TABLE 10 - Correlation Matrix for Arterial and Mixed Venous Blood Gas and Acid-Base Variables of Young Swine
- TABLE 11 - Correlation of Femoral Artery Blood Gas and Acid-Base Variables to Ventilatory Variables of Young Swine

APPENDIX

TABLE 1  
Animal Characteristics and Experimental Conditions at  
the Time of Blood Gas and Acid-Base Measurement

Characteristic	Mean	Range	S.D.	SEM
Body Weight (kg)	29.7	22.7-40.0	5.92	1.53
Hemoglobin (g/dl)	12.3	11.0-13.1	0.73	0.19
Inspired O <sub>2</sub> (%)	38	30-50	7.6	3.1
Insp. Min. Vol (l/min, BTPS)	6.5	3.5-8.4	1.38	0.36
Resp Freq. (breaths/min)	15	11-18	1.8	0.5
Tidal Vol. (ml/BTPS)	426	320-550	72	19

TABLE 2

Arterial and Venous P O<sub>2</sub> Values\* for Young Swine

Vessel	Mean	Range	S.D.	S.E.M.
Femoral Artery	97	86-115	8.6	2.2
Pulmonary Artery	36	26- 44	4.2	1.1
Ant. Vena Cava	39	33- 47	4.2	1.1
Post. Vena Cava	35	26- 42	4.4	1.1
Int. Jugular Vein	39	29- 51	6.4	1.7
Femoral Vein	35	25- 44	4.7	1.2
Coronary Sinus	30	22- 35	4.8	1.2

\* All values expressed as torr

TABLE 3

Arterial and Venous O<sub>2</sub> Saturation Values\* for Young Swine

Vessel	Mean	Range	S.D.	S.E.M.
Femoral Artery	94	91-97	1.4	0.4
Pulmonary Artery	51	37-65	7.9	2.0
Ant. Vena Cava	58	46-67	8.1	2.1
Post. Vena Cava	49	33-63	9.3	2.4
Int Jugular Vein	52	34-72	16.0	4.1
Femoral Vein	46	32-62	14.4	3.7
Coronary Sinus	42	30-61	10.2	2.6

\* All values expressed as percent oxyhemoglobin

TABLE 4

Arterial and Venous O<sub>2</sub> Content Values\* for Young Swine

Vessel	Mean	Range	S.D.	S.E.M.
Femoral Artery	15.4	14.0-16.6	0.90	0.23
Pulmonary Artery	8.5	5.7-11.2	1.46	0.39
Ant. Vena Cava	9.5	7.1-11.4	1.60	0.41
Post. Vena Cava	8.1	6.2-10.7	1.63	0.42
Internal Jugular	9.3	5.6-12.1	1.69	0.44
Femoral Vein	8.1	5.2-10.6	1.60	0.41
Coronary Sinus	6.9	4.4- 9.0	1.51	0.39

\* All values expressed as ml O<sub>2</sub>/100 ml blood

TABLE 5

Arterial and Venous pH Values\* for Young Swine

Vessel	Mean	Range	S.D.	S.E.M.
Femoral Artery	7.399	7.373-7.415	0.0117	0.0030
Pulmonary Artery	7.335	7.306-7.408	0.0259	0.0067
Ant. Vena Cava	7.323	7.237-7.359	0.0303	0.0078
Post. Vena Cava	7.341	7.296-7.413	0.0331	0.0085
Int. Jugular Vein	7.328	7.280-7.371	0.0265	0.0069
Femoral Vein	7.302	7.232-7.343	0.0311	0.0080
Coronary Sinus	7.341	7.306-7.381	0.0243	0.0063

\* All values expressed in pH units

TABLE 6

Arterial and Venous P CO<sub>2</sub> Values\* of Young Swine

Vessel	Mean	Range	S.D.	S.E.M.
Femoral Artery	47	42-54	3.5	0.9
Pulmonary Artery	57	47-66	5.8	1.5
Ant. Vena Cava	58	49-65	5.2	1.3
Post. Vena Cava	57	46-67	5.9	1.5
Int. Jugular Vein	58	48-68	6.3	1.6
Femoral Vein	62	50-76	7.0	1.8
Coronary Sinus	57	48-67	5.3	1.4

\*All values expressed at torr

TABLE 7

Arterial and Venous Bicarbonate Values\* for Young Swine

Vessel	Mean	Range	S.D.	S.E.M.
Femoral Artery	27.6	26.0-30.0	1.87	0.48
Pulmonary Artery	29.6	26.6-33.6	2.31	0.60
Ant. Vena Cava	29.5	26.6-34.0	2.17	0.56
Post. Vena Cava	29.8	27.1-34.8	2.17	0.56
Int. Jugular Vein	29.6	26.6-33.9	2.31	0.60
Femoral Vein	29.4	26.2-34.0	2.01	0.52
Coronary Sinus	29.8	27.3-34.8	2.32	0.60

\*All values expressed as mEq/liter



TABLE 8

Arterial and Venous Base Excess Values\* for Young Swine

Vessel	Mean	Range	S.D.	S.E.M.
Femoral Artery	+2.8	+0.6 - +6.6	1.80	0.47
Pulmonary Artery	+3.9	+1.0 - +7.4	2.14	0.55
Ant. Vena Cava	+3.6	+1.0 - +7.8	2.01	0.52
Post Vena Cava	+4.3	+1.5 - +9.0	1.90	0.49
Int. Jugular Vein	+3.7	+1.4 - +7.6	2.02	0.52
Femoral Vein	+3.7	+1.1 - +7.5	2.08	0.54
Coronary Sinus	+4.7	+2.0 - +8.9	2.02	0.52

\* All values expressed as mEq/liter

TABLE 9a

Analyses of Variance: Evaluation of Differences  
in Blood Gas and Acid-Base Characteristics of  
Venous Blood Taken from Various Vascular Sites  
of Young Pigs

## ANOVA

Variable	Mean Square	F-Ratio
P O <sub>2</sub>	176.0	15.18*
Oxygen Saturation	457.6	13.33*
Oxygen Content	13.05	13.30*
pH	$3.13 \times 10^{-3}$	10.69*
P CO <sub>2</sub>	52.95	9.61*
Bicarbonate Conc	0.415	0.70
Base Excess Conc	2.400	4.76*

\*Significant difference at  $P \leq 0.05$ :  $F_{5,70} = 2.37$

TABLE 9b

Newman-Keuls Evaluation of Differences in Blood  
Gas and Acid-Base Characteristics of Venous  
Blood Taken from Various Vascular Sites of Young  
Pigs

Variable	PA	AVC	PVC	IJ	FV	CS
P O <sub>2</sub>	a	b	a	b	a	c
S O <sub>2</sub>	c	a	b	a	b	c
C O <sub>2</sub>	a	b	a	b	a	c
pH	ab	b	a	b	c	ab
P CO <sub>2</sub>	a	a	a	a	c	c
[HCO <sub>3</sub> <sup>-</sup> ]	a	a	a	a	a	a
[BE]	a	ab	c	ab	b	c

Abbreviations: PA= pulmonary artery, AVC= anterior vena  
cava, PVC= posterior vena cava, IJ= internal jugular,  
FV= femoral vein, CS= coronary sinus

Like letter in a given line indicates no significant ( $P \leq 0.05$ )  
difference between vascular sites.

TABLE 10

Correlation Matrix for Arterial and Mixed Venous Blood  
Gas and Acid-Base Variables of Young Swine

	pH	P O <sub>2</sub>	P CO <sub>2</sub>	[HCO <sub>3</sub> <sup>-</sup> ]
Femoral Artery				
P O <sub>2</sub>	-0.226			
P CO <sub>2</sub>	-0.474*	0.522*		
[HCO <sub>3</sub> <sup>-</sup> ]	-0.126	0.517*	0.871*	
[BE]	-0.002	0.582*	0.858*	0.957*
Pulmonary Artery				
P O <sub>2</sub>	0.287			
P CO <sub>2</sub>	-0.633*	-0.086		
[HCO <sub>3</sub> <sup>-</sup> ]	-0.134	0.088	0.842*	
[BE]	0.081	0.248	0.701*	0.972*

\*Significant interrelationship at  $P \leq 0.05$ :  $df=13$ ,  $r=0.441$

TABLE 11  
Correlation of Femoral Artery Blood Gas and Acid-Base  
Variables to Ventilatory Variables<sup>†</sup> of Young Swine

	F	F <sub>I</sub> O <sub>2</sub>	V' <sub>T</sub>	V' <sub>I</sub>
pH	-0.265	-0.418	0.213	-0.036
P CO <sub>2</sub>	0.457*	0.890*	-0.425	-0.078
[HCO <sub>3</sub> <sup>-</sup> ]	0.405	0.465*	0.333	-0.049
P O <sub>2</sub>	0.224	0.927*	0.015	0.207

<sup>†</sup> Ventilatory variables: f= respiratory frequency in breaths/min; F<sub>I</sub>O<sub>2</sub> = percentage O<sub>2</sub> inspired;

V'<sub>T</sub> = tidal volume in ml/kg; and V'<sub>I</sub> = inspiratory minute volume in ml/kg/min.

\*Significant interrelationships at P ≤ 0.05: df = 13,  
r = 0.441

OFFICIAL DISTRIBUTION LIST

Commander  
US Army Medical Research and Development Command  
ATTN: SGRD-SI/ Mrs. Madigan  
Fort Detrick, Frederick MD 21701

Defense Technical Information Center  
ATTN: DTIC-DDA (12 copies)  
Cameron Station  
Alexandria VA 22314

Director of Defense Research and Engineering  
ATTN: Assistant Director, Environmental and  
Life Sciences  
Washington DC 20301

The Surgeon General  
ATTN: DASG-TLO  
Washington DC 20314

HQ DA (DASG-ZXA)  
WASH DC 20310

Superintendent  
Academy of Health Sciences  
ATTN: AHS-COM  
Fort Sam Houston TX 78234

Assistant Dean  
Institute and Research Support  
Uniformed Services University of Health Sciences  
6917 Arlington Road  
Bethesda MD 20014

Commander  
US Army Environmental Hygiene Agency  
Aberdeen Proving Ground MD 21070

US Army Research Office  
ATTN: Chemical and Biological Sciences Division  
P.O. Box 1221  
Research Triangle Park NC 27709  
Biological Sciences Division  
Office of Naval Research  
Arlington VA 22217

Director of Life Sciences  
USAF Office of Scientific Research (AFSC)  
Bolling AFB  
Washington DC 20332

Director  
Walter Reed Army Institute of Research  
Washington DC 20012

Commander  
US Army Medical Research Institute of Infectious  
Diseases  
Fort Detrick, Frederick MD 21701

Commander  
US Army Research Institute of Environmental  
Medicine  
Natick MA 01760

Commander  
US Army Institute of Surgical Research  
Brooke Army Medical Center  
Fort Sam Houston TX 78234

Commander  
US Army Institute of Dental Research  
Washington DC 20012

Commander  
US Army Medical Bioengineering  
Research and Development Laboratory  
Fort Detrick, Frederick MD 21701

Commander  
US Army Aeromedical Research Laboratory  
Fort Rucker AL 36362

Commander  
US Army Biomedical Laboratory  
Aberdeen Proving Ground  
Edgewood Arsenal MD 21010

Commander  
Naval Medical Research Institute  
National Naval Medical Center  
Bethesda MD 20014

Commander  
USAF School of Aerospace Medicine  
Aerospace Medical Division  
Brooks Air Force Base TX 78235

